

Comment on “Comparative study of Higgs transition in one-component and two-component lattice superconductor models”

In their recent paper [1], Motrunich and Vishwanath report results of Monte Carlo study of the Higgs transition in two three-dimensional lattice realizations of the noncompact CP^1 model. The model—a gauge theory with two complex matter fields with $SU(2)$ invariance—is of great interest since it potentially can yield a characteristic example of the so-called deconfined critical point (DCP), provided it features a generic line of second-order Higgs transitions. The authors claim an observation of such a line in their simulations. In this comment, we question the above conclusion, and show that the data of Ref. [1], if properly processed, demonstrate the absence of scale invariance and rather suggest a generic first-order line.

A significant progress in numerically distinguishing between second-order and weak first-order transitions has been achieved recently by utilizing the flowgram method [2]. In the case when the existence of the tricritical point separating the first-order phase transition region (taking place at large enough couplings) from the second-order phase transition region (taking place at smaller couplings) is hard to reveal/rule out due to system size limitations, it is crucially important to perform the following data collapse analysis for flowgrams of quantities which are supposed to be scale invariant at the second-order phase transition. Namely, one rescales linear system size, $L \rightarrow C(g)L$, where $C(g)$ is a certain smooth and monotonically increasing function of the coupling constant $g = 1/4K$ (usually, with *a priori* known asymptotic behavior at $g \ll 1$, which in the present case is $C \propto g$ [2]). A collapse of the rescaled flows in a given interval $g \in [0, g_{\text{coll}}]$ implies that the tricritical point is either absent altogether, or at least is located *outside* the interval $g \in [0, g_{\text{coll}}]$, so that the type of the transition within the interval remains the same.

We note in passing that the flowgram method with data collapse analysis has already allowed us to arrive at a definitive conclusion about the generic first-order character of the phase transition in *both* the $U(1) \times U(1)$ DCP action [2] and $SU(2)$ -symmetric NCCP¹ model [3].

The authors of Ref. [1] do not perform the data collapse study. They *interpret* the data presented in their Fig. 13 as indicative of the tricritical point at $K \approx 0.2$ without mentioning which features in particular are supporting their conclusion. Here we perform the corresponding

length-scale renormalization analysis of the data digitized from Fig. 13 (lower panel) in Ref. [1]. We observe a convincing flow collapse with divergent flow indicating the absence of scale invariance. This behavior is inconsistent with the tricritical point at $K \approx 0.2$. The scaling function $C(K)$ is chosen to be $C(0.2) = 1$ and, then, two-parameter fit gave $C(K) = 0.0740/K + 0.00444 * (\exp(1.00/K) - 1)$. This character of the flow is strikingly similar to the one found in Refs. [2, 3]. Since the




FIG. 1: (Color online) Rescaled data from Fig. 13 in Ref. [1] demonstrating perfect collapse and diverging flow inconsistent with the scale invariance.

authors did not present flowgrams for the model NCCP II, we do not comment on the nature of the phases and transitions in it.

This work was supported by NSF under Grants Nos. PHY-0653183 and PHY-0653135.

Anatoly Kuklov¹, Munehisa Matsumoto², Nikolay Prokof'ev^{3,4}, Boris Svistunov⁴, and Matthias Troyer³

¹Department of Engineering & Physics, The College of Staten Island, CUNY, Staten Island, NY 10314.

²Department of Physics, University of California, Davis, CA 95616

³Institut für Theoretische Physik, ETH Zürich, CH-8093 Zürich.

⁴Department of Physics, University of Massachusetts, Amherst, MA 01003.

PACS numbers: 05.30.-d, 75.10.-b, 05.50.+q

[1] O.I. Motrunich and A. Vishwanath, arXiv:0805.1494.

[2] A. Kuklov, N. Prokof'ev, B. Svistunov, and M. Troyer, Ann. of Phys. **321**, 1602 (2006).

[3] A. Kuklov, M. Matsumoto, N. Prokof'ev, B. Svistunov, and M. Troyer, in preparation. Preliminary results have

been presented by A. Kuklov at the Nordita *Quantum Fluids* workshop (Stockholm, August 15 - September 30, 2007) <http://www.nordita.org/~qf2007/kuklov.pdf>; and APS March Meeting 2008 (New Orleans), abstract: S12.00006.